

# Electron Cloud Simulations: Beam Instabilities and Wake Fields

G. Rumolo and F. Zimmermann  
SL/AP, CERN, Geneva, Switzerland

## Abstract

HEADTAIL is a simulation programme developed at CERN which is aimed at studying the single-bunch instability arising from the interaction on successive turns of a single bunch with the cloud generated by the previous bunches. The code includes chromaticity, space charge tune spread, broad-band impedance and detuning with amplitude for more realistic simulation. Examples of application are shown. Transverse and longitudinal wake functions are also outputs of the HEADTAIL code.

## 1 INTRODUCTION

Two different aspects of the electron-cloud phenomenon are modelled and simulated by means of the two programmes ECLOUD [1] and HEADTAIL, developed at CERN.

The first program ECLOUD simulates the build up of the electron cloud during the passage of a bunch train. It provides information on the transverse electron distribution inside the vacuum chamber, the time evolution of the total number of electrons, the energy spectrum of electrons impinging on the wall, and the corresponding heat load [2]. These outputs have already proven extremely interesting both for comparison with existing data on the electron cloud in the SPS [3, 4, 5] and for extrapolation to the LHC, like the estimation of heat load [6] or of the pressure rise in the experimental areas [4]. Furthermore, the electron density value at saturation can be used as an input to study the possible destabilizing effect of the cloud on a bunch that goes through it.

The second program HEADTAIL models the interaction of a single bunch with an electron cloud on successive turns. The cloud is assumed to be generated by the preceding bunches, and is generally assumed to be initially uniform, although other initial distributions can be considered. As observed above, its density is inferred from parallel simulations with the ECLOUD code. The electrons give rise to a head-tail wake field, which amplifies any initial small deformation in the bunch shape, *e.g.*, due to the finite number of macroparticles in the simulation. Without synchrotron motion, the resulting instability resembles the beam break up in a linac [7, 8]. If synchrotron motion is included, the instability becomes similar to a regular Transverse Mode Coupling Instability (TMCI) [9]. It induces a transverse centroid motion of the longitudinal bunch slices and also a substantial emittance growth. In our simulation, the interaction between beam and cloud is calculated by computing the electric fields of either species on a two-dimensional grid, from which we then deduce the

force exerted on the macroparticles of the opposite species. The code optionally includes nonzero chromaticity in both transverse planes, the additional effect of a broad-band resonator, space charge or beam-beam, and amplitude detuning. Electrons can evolve in a field-free region or in different magnetic field configurations (strong dipole, solenoid or combined function magnet).

HEADTAIL can be used to compute the single bunch transverse wake field, the single bunch instability threshold, and the instability growth rate above the threshold. It also contains all the necessary information to extract the longitudinal wake field and the resulting potential-well distortion. Moreover, the structure of the program easily allows us to isolate single effects or to study the synergy between them. Detailed description of the code structure and applications are presented in the following sections.

## 2 DESCRIPTION OF THE SIMULATION CODE HEADTAIL

The electron cloud can act as a short-range wake field, and drive a single bunch instability [10, 11]. Such kind of instability is held to be responsible for the vertical emittance blow-up that is observed at the KEK B factory [12], and also for that in SPS. Next we describe the model that we have employed to simulate single bunch effects due to the electron cloud.




Figure 1: Schematic of the simulation recipe.

A proton or positron bunch interacts with the electron cloud during its passage. When we study the single-bunch effects of the electron cloud, only perturbations of the cloud induced by the passing bunch are considered. All the relevant bunch and lattice parameters, as well as the average equilibrium density of the electron cloud along the ring, are basic input parameters for the simulation of the coupled motion between bunch and cloud electrons. For sim-

licity, the kick approximation is used for the action of the electron cloud on the bunch: the cloud is assumed to be localised at one or more definite positions along the ring,  $s = n \times s_{\text{el}}$  with  $n = 0, 1, \dots, (N_{\text{int}} - 1)$ . Both the cloud and the bunch are modelled as ensembles of macro-particles (with  $N_p$  bunch macro-particles and  $N_e$  macro-electrons in the cloud). The bunch is also divided into  $N_{\text{sl}}$  slices, which interact with the electron cloud after one another and cause the distortion of the initially uniform cloud distribution that can significantly affect the tail of the bunch. The principle of the simulation is illustrated in Fig. 1. The interaction between bunch particles and cloud electrons is expressed by the equations of motion:

$$\frac{d^2 \underline{x}_{p,i}(s)}{ds^2} + \underline{K}(s) \underline{x}_{p,i}(s) = \left( \frac{e}{\gamma m_p c^2} \right) \cdot \sum_{n=0}^{N_{\text{int}}-1} \underline{E}_e(\underline{x}_{p,i}(s); f_e(x, y, t)) \delta(s - n s_{\text{el}}), \quad (1)$$

$$\frac{d^2 \underline{x}_{e,j}}{dt^2} = - \left( \frac{e}{m_e} \right) (\underline{E}_p(\underline{x}_{e,j}; f_{p,\text{SL}}(x, y)) + \frac{d \underline{x}_{e,j}}{dt} \times \underline{B}_{\text{ext}}), \quad (2)$$

where the positions of electrons and bunch particles are represented by the vectors  $\underline{x}_e \equiv (x_e, y_e)$  and  $\underline{x}_p(s) \equiv (x_p, y_p, z_p)$ ,  $z = s - ct$  being a co-moving longitudinal coordinate;  $\underline{K}(s)$  is the transfer matrix with the focusing strengths between two interaction points;  $f_e(x, y)$  and  $f_{p,\text{SL}}(x, y)$  represent the distribution functions of the electron cloud and of the bunch particles contained in one slice, respectively;  $\underline{E}_{e,p}$  is the electric field of the electrons and of the beam, respectively, calculated by means of a Particle-In-Cell (PIC) algorithm adapted to our code from D. Schulte's Guinea-Pig code for beam-beam studies;  $\underline{B}_{\text{ext}}$  is an external magnetic field that can significantly influence the electron dynamics in the region where the cloud is expected to be the strongest. The field of the electron cloud acting on itself can optionally be included [13], though the effect during the bunch passage is small in most cases, and hence it is normally neglected.

The momentum changes of electrons and beam macro-particles due to their mutual attraction are computed in time steps that correspond to the different longitudinal slices into which the bunch is subdivided. Between two interactions, the bunch is propagated around the arcs of the storage ring, where the betatron motion in both planes is modelled by a rotation matrix. Synchrotron motion is also included. Hence, the beam macro-particles slowly interchange their longitudinal positions, and in particular can move across slices between turns. The effect of chromaticity is modelled by an additional rotation matrix which depends on the energy of each particle. With a further rotation, the tune shift due to space charge or beam-beam can be introduced in the bunch motion. Depending on whether this last rotation is applied around the center of the chamber or

around the center of each individual bunch slice  $\bar{x}(z)$ , it models either a beam-beam interaction or a space charge force. Amplitude detuning is optionally introduced as a tune dependence on the single particle actions,  $I_{x,y}$ . Finally, a regular transverse impedance, represented by the broad-band resonator

$$Z_{1\perp} = \frac{\omega_R}{\omega} \frac{Z_t}{1 + iQ \left( \frac{\omega_R}{\omega} - \frac{\omega}{\omega_R} \right)},$$

can optionally be included. Putting it all together, the transverse phase space coordinates of the generic bunch macro-particle are transformed over one turn according to:

$$\begin{pmatrix} x_{n+1} \\ x'_{n+1} \end{pmatrix} = \mathbf{M}_1(\delta p) \cdot \mathbf{M}_2(I_x, I_y) \cdot \left[ \mathbf{M}_{\text{sc}}(z) \begin{pmatrix} x_n - \bar{x}(z) \\ x'_n + \Delta x'_{\text{EC,Z}} - \bar{x}'(z) \end{pmatrix} + \begin{pmatrix} \bar{x}(z) \\ \bar{x}'(z) \end{pmatrix} \right].$$

Table 1: SPS parameters used in the simulations.

| variable                 | symbol         | value                         |
|--------------------------|----------------|-------------------------------|
| Circumference            | $C$            | 6900 m                        |
| Beam momentum            | $p$            | 26 GeV/c                      |
| Chamber half width       | $h_x$          | 70 mm                         |
| Chamber half height      | $h_y$          | 22.5 mm                       |
| Bunch population         | $N_b$          | $10^{11}$                     |
| Rms bunch length         | $\sigma_z$     | 30 cm                         |
| Rms energy spread        | $\delta p/p_0$ | 0.0011                        |
| Slip factor              | $\eta$         | $5.78 \times 10^{-4}$         |
| Synchrotron tune         | $Q_s$          | 0.0022                        |
| Betatron tune            | $Q_{x,y}$      | 26.6                          |
| Average beta function    | $\beta$        | 15 m                          |
| Rms hor. beam size       | $\sigma_x$     | 3 mm                          |
| Rms vert. beam size      | $\sigma_y$     | 2.3 mm                        |
| Chromaticities           | $\xi_{x,y}$    | up to 0.4                     |
| Electron-cloud density   | $\rho_e$       | $10^{12} \text{ m}^{-3}$      |
| Vertical shunt impedance | $Z_t$          | 20 M $\Omega$ /m              |
| Quality factor           | $Q$            | 1                             |
| Resonant frequency       | $\omega_R$     | $2\pi \times 1.3 \text{ GHz}$ |

### 3 SIMULATION OF THE ELECTRON CLOUD INDUCED INSTABILITY AT THE CERN SPS

As explained above, our code tracks electrons transversely along each bunch passage, and bunch particles over many turns.

The simulated electron motion reveals how electrons are progressively focused towards the bunch region as the bunch goes by. In Fig. 2 the electron phase spaces and distributions are shown, as they appear at the end of the interaction with one bunch (having started from transverse uniform distributions). The simulation has been performed

here with the PIC code; the agreement with the evolution predicted by a soft-Gaussian approximation is excellent, as may be deduced from the pictures in Ref. [14].

As far as SPS simulations are concerned (for parameters,




Figure 2: Horizontal and vertical phase space of the electrons after a bunch has gone through (top pictures), and relative distributions (bottom pictures).

see Table 1), It is interesting to observe in Fig. 3 how an SPS bunch containing  $N_b = 8 \times 10^{10}$  protons would suffer a strong dipole mode instability under the effect of the broad-band impedance alone, but now this effect gets actually damped by space charge and enhanced by the electron cloud. The instability manifests itself only in the vertical plane. There are at least two good reasons to account for that: firstly, in the vertical plane the impedance is larger [15], and secondly, there is evidence that in the SPS the electron cloud is mostly localised in the arcs, where there is a strong vertical magnetic field and therefore the electrons are mostly pinched vertically by the passing bunch. A positive chromaticity can strongly damp the instability, as shown in Fig.4 where the emittance growth over 12 ms is plotted for three different values of chromaticity.

In the SPS at 26 GeV/c space charge seems to play a




Figure 3: Evolution of the centroid vertical position of an SPS bunch over 500 turns for the three labelled cases.




Figure 4: Vertical emittance versus time for three different values of chromaticity. Broad band impedance [15] and space charge are included.

key role, since it actually causes a coherent centroid motion even where electron cloud alone would cause only incoherent emittance growth. Comparison of Figs. 5 and 6 reveals that space charge renders the beam motion more unstable and violent. In particular, it leads to slice centroid oscillations along the bunch. On the other hand, the simulation without space charge shows only a persistent emittance growth, occurring more or less uniformly along the bunch. The different signatures of the simulated instability for these two cases resemble the difference between the actual beam observations at SPS and KEKB. At the SPS injection momentum of 26 GeV/c, the beam is still affected by space charge forces, which according to our simulations might be a reason for the observed difference.




Figure 5: Simulated vertical bunch shape (centroid and rms beam size) after 0, 250, and 500 turns in the CERN SPS without proton space charge.




Figure 6: Simulated vertical bunch shape (centroid and rms beam size) after 0, 250, and 500 turns in the CERN SPS with proton space charge at 26 GeV/c.

#### 4 TRANSVERSE AND LONGITUDINAL WAKE FIELDS IN THE CERN SPS

Also wake fields can be calculated by using the HEADTAIL code. In the simulation, we displace one bunch slice (for instance, vertically by an amount  $\Delta y \propto \sigma_y$ ), and then we evaluate the electron cloud response in terms of electric field on axis ( $x = y = 0$ ). Normalizing this field by the amount of displacement and the number of particles contained in the displaced slice, we obtain the dipole wake function on axis (in  $\Omega s^{-1} m^{-1}$ , after multiplication by the factor  $m\gamma c^2/e^2$ ). As the field on axis is not directly related to the force exerted by the cloud on the slices that follow the displaced one, we can also evaluate an averaged dipole wake function from the net force caused by a displaced slice on later portions of the beam. In this case, instead of looking only at the field on axis, we calculate the overall force exerted by the distorted cloud on all the particles contained in one slice, and then divide by the total charge in that slice to obtain an effective electric field. Shapes in the two cases appear quite different, as shown in Figs. 7 and 8. Note that the two definitions of the wake would yield the same result for a conventional dipole wake field. Wake functions on axis reach much larger values and exhibit a spiky structure that is smoothed out to a more regular profile when the integration over the bunch slice is carried out. These plots correspond to an almost round beam in an SPS field-free region and are calculated for a longitudinally uniform bunch distribution. In a dipole region, the horizontal wake tends to disappear, and the vertical one becomes also weaker (see Fig. 9). Fig. 10 shows the averaged dipole wake function for offset first slice and a Gaussian bunch distribution in a dipole field region.

Even though the kick approximation allows us to use a two-dimensional model to study transverse effects, the electron cloud is in reality distributed more or less uniformly around the ring, and thus generates a longitudinal wake field which may give rise to potential well distortion and eventually micro-wave instability. The longitu-




Figure 7: Horizontal and vertical averaged dipole wake functions for a uniform SPS bunch, evaluated displacing three different bunch slices at  $t = 0, 3/10\Delta t_b, 3/5\Delta t_b$ . The simulation has been carried out in field-free region.




Figure 8: Horizontal and vertical wake functions on axis for a uniform SPS bunch, evaluated displacing three different bunch slices at  $t = 0, 3/10\Delta t_b, 3/5\Delta t_b$ . The simulation has been carried out in field-free region.

dinal field arises primarily from the accumulation of electrons near the center of the bunch during its passage. The code HEADTAIL does not directly compute the longitudinal electric field. However, we can obtain the latter from the time evolution of the transverse electron distribution. The idea consists in reconstructing the 3-dimensional electron distribution by identifying the time during the bunch passage with the longitudinal position along the bunch. This distribution can then be post-processed in order to compute the longitudinal electric field on a 3-dimensional grid. We assume that the initial electron distribution is unperturbed and uniform, and assign a homogeneous charge distribution to the region of the grid which lies in front of the bunch. The electric field is calculated on the 3-D grid points using a cloud-in-cell algorithm, and is then multiplied by the factor  $\Delta z/C$  to account for the fact that the real electrons are distributed all around the circumference  $C$ . Figure 11 displays the longitudinal electric field  $E_z(z)$  due to the electron cloud simulated for a Gaussian bunch

in the SPS. The bunch profile is also indicated. The field is negligibly small (in agreement with the estimation from a full 3-D plasma physics code [16]), less than 10 V/m. To estimate the possible bunch distortion due to this field, we assume a Gaussian energy distribution and compute the longitudinal bunch profile expected for the electron-cloud potential well using

$$\rho(z) = \rho(0) \exp \left[ -\frac{1}{2} \left( \frac{\omega_s z}{\eta c \sigma_\delta} \right)^2 \right] - \frac{r_0}{\eta \sigma_\delta^2 \gamma C} \int_0^z dz' \int_{-\infty}^{z'} dz'' \rho(z'') W_0(z' - z'') \Bigg] = \rho(0) \exp \left[ -\frac{1}{2} \left( \frac{\omega_s z}{\eta c \sigma_\delta} \right)^2 - \frac{r_0}{\eta \sigma_\delta^2 \gamma C} \int_0^z dz' W(z') \right], \quad (3)$$

where the longitudinal wake  $W(z)$ , wake function from a Gaussian bunch, is related to the longitudinal electric field estimated from the HEADTAIL code by

$$W(z) \approx \frac{E_z(z)}{e} \left( \frac{4\pi}{Z_0 c} \right) C.$$

We like to call Eq. (3) the *quasi-Haissinski solution*. Un-




Figure 9: Horizontal and vertical averaged dipole wake functions for a uniform SPS bunch, evaluated displacing the first bunch slice, in a dipole field region.

like the real Haissinski equation [18] for an ordinary wake field, Eq. (3) is not self-consistent, since the field  $E_z(z)$  on the RHS varies with the beam distribution in an unknown way. As reported in the first equality of Eq. (3), valid for a regular wake, the wake field can be expressed as a convolution between the distribution function and the Green function wake  $W_0$ : the equation can then be solved numerically for  $\rho(z)$ . For the electron cloud case, such Green function  $W_0$  is not known, and it may not even exist owing to violations of linearity and time invariance.

Nevertheless, we can use Eq. (3) to compute the bunch profile which would be formed under the influence of the additional electric field  $E_z(z)$  (neglecting its dependence




Figure 10: Horizontal and vertical averaged dipole wake functions for a Gaussian SPS bunch, evaluated displacing the first bunch slice, in a dipole field region.




Figure 11: Longitudinal electric field due to the electron cloud for a Gaussian bunch in the SPS. Bunch head is on the left.

on the bunch profile itself), and compare this with the initial distribution. A discrepancy would indicate a significant potential-well distortion, and therefore would require a few more iterations to determine the self-consistent bunch profile.

However, Fig. 12 shows that the initial and predicted distributions are very similar, and hence we do not expect a large effect of the electron cloud on the longitudinal bunch shape in the SPS. We note that the modified distribution is shifted slightly forward, which compensates for the additional energy loss due to the cloud.

## 5 APPLICATION TO THE KEKB

So far we have shown results for the CERN SPS ring. It is interesting to show, as further applications of the HEADTAIL code, a few results from simulations of the KEKB Factory Low Energy Ring. Parameters that we have used in our simulations are summarized in Table 2.

Figures 13 and 14 show horizontal and vertical centroid




Figure 12: Equilibrium bunch density computed from the wake for a Gaussian bunch in the SPS. The Gaussian is slightly shifted.




Figure 13: Horizontal centroid motion of a KEK bunch interacting with an electron cloud. Chromaticity  $\xi_x$  was set to zero. No unstable motion is visible.

Table 2: Simulation parameters for KEKB LER.

| variable               | symbol           | value                    |
|------------------------|------------------|--------------------------|
| Circumference          | $C$              | 3016 m                   |
| Relativistic factor    | $\gamma$         | 6850                     |
| Chamber radius         | $b$              | 47 mm                    |
| Bunch population       | $N_b$            | $3.3 \times 10^{10} e^+$ |
| Bunch spacing          | $T_{\text{sep}}$ | 8 ns                     |
| Rms bunch length       | $\sigma_z$       | 4 mm                     |
| Rms hor. beam size     | $\sigma_x$       | 420 $\mu\text{m}$        |
| Rms vert. beam size    | $\sigma_y$       | 60 $\mu\text{m}$         |
| Transverse tunes       | $Q_{x,y}$        | 45.53/44.08              |
| Synchrotron tune       | $Q_s$            | 0.015                    |
| Slippage factor        | $\eta$           | $1.8 \times 10^{-4}$     |
| Average beta function  | $\beta_y$        | 10 m                     |
| Chromaticities         | $\xi_{x,y}$      | up to 0.35               |
| Solenoid field         | $B_z$            | 30 G                     |
| Electron cloud density | $\rho_e$         | $10^{12} \text{ m}^{-3}$ |




Figure 14: Vertical centroid motion of a KEK bunch interacting with an electron cloud. Chromaticity  $\xi_y$  was set to zero. The coherent dipole oscillation reveals an instability.

motion for a nominal bunch under the action of an electron cloud with density  $10^{12} \text{ m}^{-3}$  and supposed uniformly distributed all along the ring. Chromaticity was set to zero in this case. A dipole motion is observable only in the vertical direction, probably because of the flatness of the KEK bunch. To illustrate the beneficial effect of chromaticity on the instability, we show first in Fig. 15 how the vertical centroid motion is damped for high vertical chromaticity. Then, in Fig. 16 the emittance growths relative to different chromaticity values are plotted on the same graph. It is clear that higher positive chromaticities can damp the coherent emittance growth otherwise predicted for zero or very low chromaticity. For the purpose of code benchmarking and validation, we show in Fig. 17 the same chromaticity scan for KEK done by K. Ohmi [17] with his PEI code. The agreement with our results is excellent.

The influence of a solenoid field on the electron cloud driven single bunch instability can also be of interest, and it has been studied in connection with the the KEK too, since there are solenoids along 90% of the ring. In Figs. 18

and 19 the horizontal and vertical wake functions are plotted for a KEK nominal bunch and the nominal value of the solenoid field ( $B_{\text{sol}} = 3 \text{ mT}$ ). The difference between the two figures is that the horizontal and vertical displacements were separately applied to the first bunch slice: nevertheless, the expected effect of exciting a wake in the orthogonal plane is in neither case quite visible in both planes in spite of the coupling in the electron motion due to the solenoid. This is because the KEK bunch is very short and the solenoid field not strong enough to efficiently couple planes over one bunch pass (electron cyclotron period for  $B_{\text{sol}} = 3 \text{ mT}$  is  $T_c = 12 \text{ ns}$ , and the KEK bunch length is  $\Delta t_b = 0.053 \text{ ns}$ ). In Ref. [13] it was shown how for a long SPS bunch in a solenoid a strong effect was visible in both planes even after displacing a bunch slice only in one plane. When the coupling is effective, the wake field becomes much weaker and the instability is likely to be cured [13], as appears from the loss of pinching shown in Fig. 20 for an SPS bunch propagated through an electron cloud in a solenoid field region ( $B_z = 0, 2.5, 10 \text{ mT}$ ).






Figure 15: Vertical centroid motion of a KEK bunch interacting with an electron cloud. Chromaticity  $\xi_y$  was set to 0.35. The coherent dipole oscillation is damped with respect to the case with zero chromaticity.




Figure 16: Vertical rms-size growth of a KEK bunch over 500 turns for different values of chromaticity (labelled in  $Q'_y = \xi_y \cdot Q_y$ ). Chromaticity helps against the e-cloud instability. All plots have been obtained using the HEADTAIL code.

## 6 CONCLUSIONS AND OUTLOOK

In conclusion, in this paper we have described the code HEADTAIL that was developed at CERN in order to study the degrading effect of an electron cloud on the single bunch. Examples of application have been shown, with special emphasis to the CERN SPS. Growth rate of the instability and beneficial effect of chromaticity, as experimentally observed [19], have been reproduced by simulations including electron cloud, space charge and broad band impedance. It has also been highlighted that space charge plays a key role in destabilizing the centroid beam motion: when its effect is neglected only an incoherent emittance growth is predicted to occur. Transverse and longitudinal wake functions have been calculated by means of the HEADTAIL code. Transverse wakes can be of great interest to predict the instability threshold from the TMCI theory, although this theory




Figure 17: Vertical rms-size growth of a KEK bunch for different values of the chromaticity (labelled in  $Q' = \xi_y \cdot Q_y$ ). Chromaticity helps against the e-cloud instability. All plots have been obtained using the PEI code by K. Ohmi (Courtesy K. Ohmi).




Figure 18: Horizontal and vertical wake functions along a KEK bunch that goes through an electron cloud inside a 30 G solenoid field region. The bunch head was displaced only horizontally by an amount  $\Delta x = \sigma_x/10$ .

can only be applied in first approximation, and needs to be adapted to this particular case. The electron cloud wake fields are not conventional, because they strongly depend on the position along the bunch from where they get excited. Work is being carried out to take into account this feature in the theory [20]. Assuming the electron distributions at subsequent times to be those at subsequent longitudinal positions (with the appropriate scaling factor), the longitudinal wake field has also been evaluated from the output of the HEADTAIL code, and found to be small. It can only slightly affect the bunch shape and is not likely to be responsible for any microwave instability. The results of the HEADTAIL code have been benchmarked against the predictions of Ohmi's PEI code: the beneficial effect of chromaticity on the electron cloud single bunch instability for a KEK bunch has been reproduced by both codes in excellent agreement.




Figure 19: Horizontal and vertical wake functions along a KEK bunch that goes through an electron cloud inside a 30 G solenoid field region. The bunch head was displaced only vertically by an amount  $\Delta y = \sigma_y/10$ .




Figure 20: Electron densities along the x-axis after 4/5 of the SPS bunch has passed through the cloud and for  $B_z = 0, 2.5, 10$  mT. This simulation result shows how the pinching effect disappears when the solenoid field is strong enough as to significantly couple the electron motion over one bunch pass.

## 7 ACKNOWLEDGEMENTS

The authors would like to thank F. Ruggiero, D. Schulte, G. Arduini, K. Cornelis, E. Métral, M. Giovannozzi, A. Rossi, K. Ohmi, E. Perevedentsev, and T. Katsouleas for helpful discussion and information.

## 8 REFERENCES

- [1] F. Zimmermann and G. Rumolo, “Electron Cloud Simulation: Build Up and Related Effects”, these proceedings
- [2] G. Rumolo and F. Zimmermann, “Practical User Guide for ECLOUD” CERN-SL-Note-2002-016 (AP)
- [3] V. Baglin and B. Jenninger, “SPS electron cloud heat load measurement with WAMPAC and simulation”, these proceedings
- [4] A. Rossi, G. Rumolo and F. Zimmermann, “Simulation Study of Electron Cloud in the Experimental Regions of the LHC”, these proceedings
- [5] M. Jimenez *et al.*, “Electron-Cloud Observations in the SPS”, these proceedings
- [6] F. Zimmermann, “Electron Cloud in the LHC”, these proceedings
- [7] K. Ohmi and F. Zimmermann, Phys. Rev. Lett. **85**, 3821 (2000).
- [8] G. Rumolo, F. Ruggiero and F. Zimmermann, Phys. Rev. ST Accel. Beams, **4**, 012801 (2001).
- [9] K. Ohmi, F. Zimmermann and E. Perevedentsev, Phys. Rev. E **65** (2002) 016502
- [10] F. Zimmermann, “Electron-Cloud Studies for the Low-Energy Ring of KEKB,” CERN SL-Note-2000-004 AP (2000).
- [11] K. Ohmi and F. Zimmermann, “Head-Tail Instability caused by Electron Cloud in Positron Storage Rings”, CERN-SL-Note-2000-015 AP (2000).
- [12] H. Fukuma, *et al.*, “Observation of Vertical Beam Blow-Up in KEK Low Energy Ring”, in Proc. of EPAC 2000, Vienna (2000).
- [13] G. Rumolo and F. Zimmermann, “Contributions of the SL-AP Group to the Two-stream Instabilities Workshop”, CERN SL-2001-067 (2001).
- [14] G. Rumolo and F. Zimmermann, “Theory and Simulation of the Electron Cloud Instability” Chamonix XI, CERN SL-2001-003 (DI)
- [15] H. Burkhardt, G. Rumolo and F. Zimmermann, “Measurements of SPS Single Bunch Coherent Tune Shifts and Head-Tail Growth Rates in the Year 2001” CERN, SL-Note-2001-043 (MD)
- [16] T. Katsouleas *et al.*, “Plasma Modelling of Collective Wakefields in Electron Clouds”, these proceedings.
- [17] K. Ohmi, “Particle-In-Cell Simulation of Beam-Electron Cloud Interactions” Proc. PAC 2001, Chicago, USA (2001).
- [18] J. Haissiniski, Nuovo Cimento **18B**, 72 (1973)
- [19] K. Cornelis *et al.*, “Electron Cloud Instability in the SPS”, these proceedings.
- [20] E. Perevedentsev *et al.*, “Head-Tail Instability Caused by Electron-Cloud”, these proceedings.